

Discovery of a Large-scale Wall in the Direction of Abell 22

Kevin A. Pimbblet,^{1,2} Alastair C. Edge,³ Warrick J. Couch⁴

¹ *pimbblet@physics.uq.edu.au*

² *Department of Physics, University of Queensland, Brisbane, QLD 4072, Australia*

³ *Department of Physics, University of Durham, South Road, Durham, DH1 3LE, UK*

⁴ *School of Physics, University of New South Wales, Sydney, NSW 2052, Australia*

DRAFT: 2 FEBRUARY 2008 — DO NOT DISTRIBUTE

ABSTRACT

We report on the discovery of a large-scale wall in the direction of Abell 22. Using photometric and spectroscopic data from the Las Campanas Observatory and Anglo–Australian Telescope Rich Cluster Survey, Abell 22 is found to exhibit a highly unusual and striking redshift distribution. We show that Abell 22 exhibits a foreground wall-like structure by examining the galaxy distributions in both redshift space and on the colour-magnitude plane. A search for other galaxies and clusters in the nearby region using the 2dF Galaxy Redshift Survey database suggests that the wall-like structure is a significant large-scale, non-virialized filament which runs between two other Abell clusters either side of Abell 22. The filament stretches over at least $> 40 \text{ h}^{-1} \text{ Mpc}$ in length and $10 \text{ h}^{-1} \text{ Mpc}$ in width at the redshift of Abell 22.

Key words: galaxies: clusters: individual: Abell 22 – surveys – large-scale structure of Universe

1 INTRODUCTION

Hierarchical structure formation modelling (Zeldovich, Einasto & Shandarin 1982; Katz et al. 1996; Bond, Kofmann & Pogosyan 1996; Jenkins et al. 1998; Colberg et al. 2000) succinctly demonstrates how large scale structure such as galaxy clusters grow from the non-isotropic infall and accretion of smaller components. Material infalling toward galaxy clusters is found to be preferentially situated along galaxy filaments (Bond, Kofmann & Pogosyan 1996). These filaments are, in turn, observed to stretch between galaxy clusters at all redshifts (e.g. Ebeling, Barrett, & Donovan 2004; Pimbblet & Drinkwater 2004; Colberg, Krughoff & Connolly 2004; Pimbblet, Drinkwater & Hawkrigg 2004; Dietrich et al. 2004; Gal & Lubin 2004; Durret et al. 2003; Scharf et al. 2000; Doroshkevich et al. 2000; Colberg et al. 1999; Geller et al. 1997; Bond, Kofman & Pogosyan 1996; Shectman et al. 1996; Geller & Huchra 1989) and are also observed near very isolated clusters (Durret et al. 2003; Kodama et al. 2001). Since galaxy filaments can comprise over 40 per cent of the total cluster mass at clustocentric radii of $4\text{--}6.5 \text{ h}^{-1} \text{ Mpc}$ (Colberg et al. 1999; see also Cen & Ostriker 1999) their direct observation remains paramount to our understanding of cluster growth.

Significantly, not all galaxy filaments are the same. They exist in many morphologies and varied lengths (Colberg, Krughoff & Connolly 2004; CKC herein; Pimbblet, Drinkwater & Hawkrigg 2004; PDH herein). Shorter filaments are generally found between close cluster pairs;

longer filaments are much rarer (e.g. straight filaments of $30 \text{ h}^{-1} \text{ Mpc}^*$ in length are at least 5 times as rare as those of $10 \text{ h}^{-1} \text{ Mpc}$ in length; PDH). In the nomenclature of PDH, Type III inter-cluster filament morphologies (‘walls’ or ‘sheets’ of galaxies) are exceedingly rare, comprising *at most* no more than 3 per cent of the total filament population (CKC; PDH). In this work, we report on the discovery of a large-scale ($> 40 \text{ h}^{-1} \text{ Mpc}$) wall-like filament in the direction of Abell 22 (X-ray centre is at $\alpha = 00^{\text{h}} 20^{\text{m}} 38^{\text{s}}$, $\delta = -25^{\circ} 43' 19''$; $cz = 42676 \pm 98 \text{ kms}^{-1}$; Pimbblet 2001) found in the Las Campanas Observatory and Anglo–Australian Telescope Rich Cluster Survey (LARCS; e.g. Pimbblet et al. 2001). Briefly, LARCS is a long-term project to study a statistically-reliable sample of 21 of the most luminous X-ray clusters at intermediate redshifts ($0.07 < z < 0.16$) in the southern hemisphere. We are mapping the photometric, spectroscopic and dynamical properties of galaxies in rich cluster environments at $z \sim 0.1$, tracing the variation in these properties from the high-density cluster cores out into the surrounding low-density field beyond the turn-around radius. For the most massive clusters at $z \sim 0.1$, the turn-around radius corresponds to roughly 1 degree or a $10 \text{ h}^{-1} \text{ Mpc}$ radius (O’Hely et al. 1998; Pimbblet 2001) and therefore we have obtained panoramic CCD imaging covering 2-degree diameter fields, as well as spectroscopic coverage of

* We use $H_0 = 100 \text{ h km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$ throughout this work. Further, all quoted coordinates are J2000 compliant.




Figure 1. Velocity histogram for Abell 22 with inset showing a magnification of the central region. The bins are 1000 km s^{-1} wide in the main plot and 400 km s^{-1} wide in the inset panel. The downward arrow denotes the velocity of Abell 22 as computed by Pimblet (2001). There is a lot of significant structure in the foreground of Abell 22.

these fields (e.g. Pimblet et al. 2001; O’Hely 2000). The imaging comes from *B* and *R*-band mosaics taken with the 1-m Swope telescope at Las Campanas Observatory, while the spectroscopy comes from the subsequent follow-up with the 400-fibre 2dF multi-object spectrograph on the 3.9-m Anglo-Australian Telescope (AAT).

The plan of this paper is as follows: In §2 we examine the unusual velocity structure of Abell 22 and the foreground wall-like structure. We then divide our observations into redshift bins and examine the distribution of the constituent galaxies on the colour-magnitude plane in §3. In §4 we present a search for other local clusters to determine if the wall-like structure is a part of a larger-scale object. We summarize our findings in §5.

2 VELOCITY AND SPATIAL STRUCTURE

The photometric data reduction for LARCS is presented in Pimblet et al. (2001) whilst the spectroscopic reduction will be presented in Pimblet et al. (in prep.).

Observations of Abell 22 using 2dF (Pimblet 2001) have yielded redshifts for 345 galaxies. From these, it is possible to see the presence of an unusual structure from the redshift distribution, shown in Figure 1. Figure 2 shows the position of the galaxies along the line of sight with respect to their angular position on the sky.

Abell 22 has a central peak which coincides with *some* previously published estimates of the clusters redshift (e.g. Dalton et al. 1994). The peak of the distribution in the velocity histogram is unusually broad and contains multiple sub-




Figure 2. Wedge plots of right ascension and declination versus redshift in the direction of Abell 22. The X-ray centre of the cluster is located at $\alpha = \delta = 0$. Note the ensemble of galaxies in the foreground of the cluster. Although one may expect this plot to exhibit a ‘finger of God’ effect for the cluster, arising from the distorting effects in redshift space, this is not seen as the elongation in redshift space of the cluster is small in comparison to the depth of redshift space covered.

peaks. The wedge plot (Figure 2) reveals that Abell 22 is a highly complex structure. The foreground galaxies are striking: a non-virialized wall-like structure appears to stretch across the entire RA range of the observations, but is more limited in scope (and offset from the cluster) in Dec. It is little wonder, therefore, that other redshift estimates for Abell 22 differ from that of Dalton et al. (1994) by as much as $\sim 3000 \text{ km s}^{-1}$ (e.g. Struble & Rood, 1999).

3 THE COLOUR-MAGNITUDE RELATION

We now split our sample up into seven equal redshift bins from 31875 km s^{-1} to 45000 km s^{-1} and construct colour-magnitude diagrams for each bin. These diagrams are displayed in Figure 3.

Table 1. Fraction of blue galaxies, f_B , within each redshift bin; where f_B is defined here as being the fraction brighter than $M^* + 1$ possessing a rest frame colour $\Delta(B - V) = -0.2$ bluer than the fitted CMR of Pimbblet et al. (2002) over all galaxies brighter than $M^* + 1$. The column headed $N(M < M^* + 1)$ is the number of galaxies brighter than $M^* + 1$.

Redshift Bin (kms^{-1})	$N(M < M^* + 1)$	Blue Fraction, f_B
43126-45000	19	0.05 ± 0.05
41251-43125	33	0.18 ± 0.07
39376-41250	9	0.22 ± 0.16
37501-39375	21	0.29 ± 0.12
35626-37500	25	0.52 ± 0.14
33751-35625	32	0.31 ± 0.10
31876-33750	26	0.35 ± 0.12




Figure 3. Colour-magnitude diagrams for each of the seven redshift bins. The vertical dashed line denotes the $M^* + 1$ magnitude division used during the 2dF observations (Pimbblet 2001). The dotted lines are the 1σ error to the inner ($r_p < 1\text{Mpc}$) region CMR fit from Pimbblet et al. (2002).

Pimbblet et al. (2002) investigated the environmental dependence of galaxy colours within a sample of eleven LARCS clusters; including Abell 22. They construct colour-magnitude diagrams divided into radial intervals for these clusters and fitted the colour-magnitude relation (CMR; e.g. Visvanathan & Sandage 1977) using a biweight method. Here, we use the errors on their CMR fit to Abell 22 as a *guideline* for colour magnitude diagrams constructed for each of the redshift bins (Figure 3).

The CMR distribution for the redshift bins nearer to the filament are qualitatively similar to those nearer to the cluster (Figure 3). To quantify the similarity of the distributions, we derive the blue fraction, f_B , of galaxies in each redshift bin. Here, we define f_B to be the fraction of galaxies brighter than $M^* + 1$ possessing a rest frame colour that is $\Delta(B - V) = -0.2$ bluer than the fitted CMR of Pimbblet et al. (2002). This differs from the classical Butcher-Oemler (1984; BO; see Pimbblet 2003 for a review) definition of f_B (i.e. our $M^* + 1$ magnitude cut does not correspond to the $M_V = -20$ magnitude cut used by BO and we have not applied a limiting clustocentric radius), but this is a sound approach as we are simply interested in internal differences in f_B between our redshift bins. Table 1 lists the blue fractions for the redshift bins together with associated Poissonian errors. The blue fraction increases from the upper two redshift bins (i.e. Abell 22) to the lower redshift bin (i.e. the wall-like structure), but only by $1-2\sigma$. Given the large errors on these numbers and the low number of galaxies in some bins (see Table 1) we do not view this as a highly significant change. Moreover, the blue fraction of the filament is very comparable to that found by Pimbblet & Drinkwater (2004).

4 DISCUSSION

To determine whether the wall-like structure displayed in Figure 2 may be a part of a more extended object we utilize the NASA/IPAC Extragalactic Database (NED) to look for any near-by (within 3 degrees on the sky) clusters of galaxies at similar redshifts. Four clusters in addition to Abell 22 are found and we list all five of these in Table 2. Given the redshifts of Abell 14 and APMCC 029, it is unlikely that they are associated with the filament. Meanwhile, Abell 22 and Abell 47 certainly reside in the background of the filament and are likely unconnected due to the huge physical scale between them and the filament. We note that Abell 15, however, resides at a redshift that is similar to the filament, but is Abell 15 connected to the filament?

Since this region overlaps (Pimbblet et al. 2001) with the 2dF Galaxy Redshift Survey (2dFGRS; Colless et al. 2001) we now use the final release of the 2dFGRS (see <http://www.mso.anu.edu.au/2dFGRS/>) to search for any galaxies with measured redshifts in the range 33751–35625 kms^{-1} , thereby encompassing our final redshift bin. The result of both of these searches are illustrated in Figure 4. We note that the 2dFGRS final data release does not extend above $\delta \sim -24.9$ but does encompass Abell 15, much of Abell 22 and partial amounts of Abell 47 (Figure 4). We can therefore use 2dFGRS to try to answer the question of whether Abell 15 is connected to the filament. Since the velocity dispersion for Abell 15 is $\sigma_z \sim 500 \text{ kms}^{-1}$ † (De Propris et al. 2002), the filament easily resides within its infall regions (Diaferio & Geller 1997) at $2 < \Delta v / \sigma_z < 4$ (see Carlberg, Yee & Ellingson 1997). An inspection of galaxy

† De Propris et al. (2002) note that their observations for Abell 15 are 54 per cent complete ($N_{gal} = 24$), hence we use a velocity dispersion of $\sim 500 \text{ kms}^{-1}$ as an order of magnitude estimate only. The actual value calculated by them is $\sigma_z = 459_{-82}^{+104} \text{ kms}^{-1}$.

Table 2. Position and redshifts of all clusters with measured redshifts within three degrees of Abell 22 found using NED. The column headed ‘Redshift Reference’ gives the source of the quoted redshift.

Cluster Name	RA (J2000)	Dec	Distance from Abell 22 (′)	Redshift (kms ⁻¹)	Redshift Reference
Abell 22 [‡]	00 20 34.8	-25 41 51	n/a	42658	Pimblet (2001)
Abell 15	00 15 13.7	-26 01 19	75	36043	De Propriis et al. (2002)
APMCC 029	00 15 04.0	-24 00 04	126	18887	Dalton et al. (1997)
Abell 14	00 15 13.9	-23 53 19	131	19636	Struble & Rood (1999)
Abell 47	00 30 35.9	-24 09 26	165	41431	Struble & Rood (1999)

[‡] Coordinates for Abell 22 presented here are from NED. They differ slightly to the coordinates of the cluster’s X-ray centre as noted in section 1.

surface density contours around Abell 15 and Abell 47 also shows that these clusters are elongated in the direction of the filament (Figure 4; see Binggeli 1982; Plionis et al. 2003 and references therein) meaning they are very likely to be connected.

We suggest that the wall-like structure is a significant non-virialized overdensity in galaxies which runs from (at least) Abell 15, across Abell 22 and toward Abell 47. If so, then the filament stretches over at least $> 40 h^{-1}$ Mpc at this redshift. The surface density (computed in the same way as Pimblet et al. 2002) of the filament is typically ~ 10 galaxies per square h^{-1} Mpc. Such a surface density is very similar to that found in other filaments (Ebeling, Barrett & Donovan 2004; PDH). As the filament connects two clusters, we cannot yet consider it to be of the same ilk as the Great Wall noted by Geller & Huchra (1989). It may however, be taken as evidence that the region around Abell 22 is a supercluster candidate.

5 SUMMARY

We have presented evidence for a new large-scale wall in the direction of Abell 22.

- The velocity structure of Abell 22 is highly unusual: the cluster appears to extend in redshift space and merges with foreground galaxy structure. These foreground galaxies are highly striking: a wall-like structure appears to stretch across the entire RA range of the observations. It is probable that the presence of this wall-like structure has caused previous estimates of Abell 22’s redshift to differ significantly.
- Colour-magnitude diagrams for the redshift bins are qualitatively similar and the blue fraction of galaxies does not change significantly between the cluster and the foreground filament.
- The wall appears to be part of a large-scale filament that passes in front of Abell 22 and Abell 47, and is likely connected to Abell 15, which is at a similar redshift to the filament and elongated toward it. At this redshift, the filament stretches over at least $40 h^{-1}$ Mpc. We suggest that this region is a supercluster candidate.

Further spectroscopic observations in the direction of Abell 47 are required to definitively determine the filaments extension toward this cluster and to find out if the structure

is part of a larger supercluster. It would also be interesting to investigate if the blue fraction and morphology of galaxies along other filaments (e.g. PDH) remained approximately constant along their length and how similar is it to clusters they connect to? Indeed the morphology-density relation (Dressler 1980) of filaments remains a relatively unstudied area.

Acknowledgements

We thank the anonymous referee for a very speedy and useful review. We also thank Mary Hawkrigg and Nathan Courtney for reading through an earlier draft of this work and providing useful comments. KAP acknowledges support from an EPSA University of Queensland Research Fellowship and a UQRSF grant. ACE acknowledges support from the Royal Society. WJC acknowledges the financial support of the Australian Research Council. We thank the Observatories of the Carnegie Institution of Washington and the Anglo-Australian Observatory for their generous support of this survey. In particular, we warmly thank Geraint Lewis and Terry Bridges for their assistance at the telescope.

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Binggeli B., 1982, *A&A*, 107, 338
Bond J. R., Kofman L., Pogossyan D., 1996, *Nature*, 380, 603
Butcher, H. & Oemler, A. 1984, *ApJ*, 285, 426 (BO)
Carlberg R. G., Yee H. K. C., Ellingson E., 1997, *ApJ*, 478, 462
Cen R., Ostriker J. P., 1999, *ApJ*, 514, 1
Colberg J. M., Krughoff K. S., Connolly A. J., 2004, *astro*, *astro-ph/0406665* (CKC)
Colberg J. M., et al., 2000, *MNRAS*, 319, 209
Colberg J. M., White S. D. M., Jenkins A., Pearce F. R., 1999, *MNRAS*, 308, 593
Colless M. et al., 2001, *MNRAS*, 328, 1039
Dalton, G. B., Maddox, S. J., Sutherland, W. J., & Efstathiou, G., 1997, *MNRAS*, 289, 263
Dalton, G. B., Efstathiou, G., Maddox, S. J., & Sutherland, W. J., 1994, *MNRAS*, 269, 151
De Propriis R., et al., 2002, *MNRAS*, 329, 87
Diaferio A., Geller M. J., 1997, *ApJ*, 481, 633




Figure 4. Spatial distribution of galaxies contained in the wall-like structure (filled triangles), galaxies contained in the 2dFGRS final data release (crosses) and all clusters with measured redshifts within 3 degrees of Abell 22 (large circles of diameter 2 degrees). Note that the 2dFGRS final data release does not extend across all of this area; its approximate extent is denoted by the dot-dashed line. The scale bar shows $10 h^{-1}$ Mpc at the central redshift of the $33751\text{--}35625 \text{ km s}^{-1}$ bin. We suggest the wall-like structure extends from beyond Abell 15, across Abell 22 and toward Abell 47.

Dietrich J. P., Schneider P., Clowe D., Romano-Diaz E., Kerp J., 2004, *astro*, *astro-ph/0406541*
Doroshkevich, A. G., Fong, R., McCracken, H. J., Ratcliffe, A., Shanks, T., & Turchaninov, V. I. 2000, *MNRAS*, 315, 767
Dressler A., 1980, *ApJ*, 236, 351
Durret F., Lima Neto G. B., Forman W., Churazov E., 2003, *A&A*, 403, L29
Ebeling H., Barrett E., Donovan D., 2004, *ApJ*, 609, L49
Gal R. R., Lubin L. M., 2004, *ApJ*, 607, L1
Geller, M. J. & Huchra, J. P. 1989, *Science*, 246, 897
Geller, M. J. et al. 1997, *AJ*, 114, 2205
Jenkins A., Frenk C. S., Pearce F. R., et al., 1998, *ApJ*, 499, 20
Katz N., Weinberg D. H., Hernquist L., Miralda-Escude J., 1996,

ApJ, 457, L57
Kodama T., Smail I., Nakata F., Okamura S., Bower R. G., 2001, *ApJ*, 562, L9
O'Hely, E., 2000, Ph.D. Thesis, University of New South Wales
O'Hely, E., Couch, W.J., Smail, I., Edge, A.C., Zabludoff, A. I., 1998, *PASA*, 15, 273
Pimbblet, K. A., 2001, Ph.D. Thesis, University of Durham
Pimbblet K. A., Smail I., Edge A. C., Couch W. J., O'Hely E., Zabludoff A. I., 2001, *MNRAS*, 327, 588
Pimbblet K. A., Smail I., Kodama T., Couch W. J., Edge A. C., Zabludoff A. I., O'Hely E., 2002, *MNRAS*, 331, 333
Pimbblet K. A., 2003, *PASA*, 20, 294
Pimbblet K. A., Drinkwater M. J., 2004, *MNRAS*, 347, 137

- Pimbblet K. A., Drinkwater M. J., Hawkrigg M. C., 2004, MNRAS, 354, L61 (PDH)
- Plionis M., Benoist C., Maurogordato S., Ferrari C., Basilakos S., 2003, ApJ, 594, 144
- Scharf C., Donahue M., Voit G. M., Rosati P., Postman M., 2000, ApJ, 528, L73
- Shethman, S. A., Landy, S. D., Oemler, A., Tucker, D. L., Lin, H., Kirshner, R. P., & Schechter, P. L. 1996, ApJ, 470, 172
- Struble, M. F. & Rood, H. J., 1999, ApJS, 125, 35
- Visvanathan N., Sandage A., 1977, ApJ, 216, 214
- Zeldovich I. B., Einasto J., Shandarin S. F., 1982, Nature, 300, 407